

POSTER PRESENTATION

Open Access

Parenteral nutrition improves nutritional status, autonomic symptoms and QoL in patients with TTR-FAP

Massimo Russo^{1*}, Gianluca Vita¹, Claudia Stancanelli², Giuseppe Vita³, Anna Mazzeo³, Sonia Messina¹

From First European Congress on Hereditary ATTR amyloidosis
Paris, France. 2-3 November 2015

Background

Transthyretin related familial amyloidotic polyneuropathy (TTR-FAP) is an inherited form of amyloidosis, leading to death in about 10 years in most cases for cardiac failure or wasting syndrome. Previous study showed that modified body mass index (mBMI) was related to time before death, duration of gastrointestinal disturbances, malabsorption and functional capacity. Furthermore, outcome after liver transplantation was greater in patients with an mBMI over 600.

Patients

We report two TTR-FAP patients, carrying respectively the Thr49Ala and the Glu89Gln mutations, in whom nutritional status worsened despite diet modification, hypercaloric supplement and two important therapeutic approaches such as liver transplant and tafamidis meglumine.

In both case, at a late stage of the disease, a peripherally inserted central catheter (PICC) was placed and the parenteral nutrition started.

Results

The parenteral nutrition added to oral nutrition allowed to improve their nutritional status and clinical conditions as documented by increase of body weight and mBMI. Moreover, we recorded reduction of autonomic symptoms including postural hypotension, nausea and diarrhoea and amelioration of quality of life.

Conclusion

Our experience suggests that parenteral nutrition administered by PICC may be useful in reducing complications

and disabilities in TTR-FAP patients, even when all dietary adjustments have been ineffective. Reasonably, the improvement in nutritional status may prolong survival in TTR-FAP patients.

Authors' details

¹Nemo Clinical Center Messina, University of Messina, 98125, Messina, Italy. ²Biomedical Department of Internal and Specialistic Medicine, University of Palermo, 90100, Palermo, Italy. ³Department of Neuroscience, University of Messina, 98125, Messina, Italy.

Published: 2 November 2015

doi:10.1186/1750-1172-10-S1-P52

Cite this article as: Russo et al.: Parenteral nutrition improves nutritional status, autonomic symptoms and QoL in patients with TTR-FAP. *Orphanet Journal of Rare Diseases* 2015 **10**(Suppl 1):P52.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Nemo Clinical Center Messina, University of Messina, 98125, Messina, Italy
Full list of author information is available at the end of the article