

RESEARCH

Open Access



A standard set of outcome measures for the comprehensive assessment of oral health and occlusion in individuals with osteogenesis imperfecta

L. Blokland^{1*}, H. Arponen^{2,3}, A. Ahmad⁴, S. Colijn^{5,6}, H. Gjørup⁷, R. John⁸, M. Li⁹, D. Mekking⁵, S. Parekh¹⁰, J. M. Retrouvey¹¹, T. Stutz Steiger¹², L. Zhou⁹ and K. Andersson^{13,14,15}

Abstract

Background Osteogenesis imperfecta (OI) is a group of inherited connective tissue disorders of varying severity characterized by bone fragility. The primary objective of this international multidisciplinary collaboration initiative was to reach a consensus for a standardized set of clinician and patient-reported outcome measures, as well as associated measuring instruments for dental care of individuals with OI, based on the aspects considered important by both experts and patients. This project is a subsequent to the Key4OI project initiated by the Care4BrittleBones foundation which aims to develop a standard set of outcome measures covering a large domain of factors affecting quality of life for people with OI. An international team of experts comprising orthodontists, pediatric dentists, oral and maxillofacial surgeons, and prosthetic dentists used a modified Delphi consensus process to select clinician-reported outcome measures (CROMs) and patient-reported outcome measures (PROMs) to evaluate oral health in individuals with OI. Important domains were identified through a literature review and by professional expertise (both CROMs and PROMs). In three focus groups of individuals with OI, important and relevant issues regarding dental health were identified. The input from the focus groups was used as the basis for the final set of outcome measures: the selected issues were attributed to relevant CROMs and, when appropriate, matched with validated questionnaires to establish the final PROMs which represented best the specific oral health-related concerns of individuals with OI.

Results Consensus was reached on selected CROMs and PROMs for a standard set of outcome measures and measuring instruments of oral health in individuals with OI.

Conclusions Our project resulted in consensus statements for standardization oral health PROMs and CROMs in individuals with OI. This outcome set can improve the standard of care by incorporating recommendations of professionals involved in dental care of individuals with OI. Further, it can facilitate research and international research co-operation. In addition, the significant contribution of the focus groups highlights the relevance of dental and oral health-related problems of individuals with OI.

Keywords Osteogenesis imperfecta, Dentinogenesis imperfecta, Malocclusion, Outcome measures, Measuring instruments

*Correspondence:

L. Blokland

l.blokland@vogellanden.nl

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Osteogenesis imperfecta (OI) is a genetic connective tissue disorder, with the characteristic symptoms of bone fragility, recurrent fractures, impaired growth, and resulting short stature [1]. Affected individuals may also present with blue sclera, joint laxity, and dentinogenesis imperfecta (DGI) [1]. OI is traditionally classified into four main types according to clinical and radiographic findings, where type I is mild with blue sclerae, type II is pre- or perinatally lethal, type III is the most severe type associated with survival of the perinatal period, and type IV is of moderate severity [2]. Autosomal dominant mutations in the *COL1A1* and *COL1A2* genes are causative in approximately 85% of cases [3–5]. More recent studies have revealed a plethora of recessive and X-linked mutations in genes related to the production and modification of collagen type I, leading to the proposal of a new OI nomenclature comprising five main types [6]. Significant efforts have also been made to adopt a dyadic naming system by systematically associate the phenotypic entity with the gene it arises from [7]. However, the classification by Sillence is still the most frequently used in clinical practice.

Dental and craniofacial aberrations are common findings in individuals with OI [8–11]. The most common oral findings in OI are DGI and malocclusion such as open bite in the anterior or posterior region of the occlusion [8, 9, 11, 12]. Teeth affected by DGI exhibit a characteristic grey-blue to brown discolouration due to dysplastic dentin [13, 14]. The enamel, which shows normal structure and mineral content, is easily fractured due to the underlying dysplastic dentin, which is prone to attrition [9]. The deciduous dentition is often more severely affected than the permanent dentition [9]. Radiographically the teeth exhibit a deviating morphology, pathognomonic for the condition, with bulbous crowns, a marked cervical constriction, pulpal obliteration, and short roots [11, 13]. DGI is classified into two subgroups based on genetic findings; DGI type I, syndromic form associated with osteogenesis imperfecta and DGI type II, non-syndromic form [14–18]. In individuals with OI, depending on the type, prevalence of DGI is estimated to be between 8 and 100% [8, 12]. More severely affected children often have a more severe dental phenotype [8, 9, 12]. Additionally, many individuals with OI present with other dental aberrations. Among these are dental agenesis, apically extended pulp chambers (taurodontism – in individuals without DGI), retained permanent second molars and abnormal craniofacial development with vertical underdevelopment of dentoalveolar structures [9, 19–23]. The oral manifestations in OI may have a significant impact on oral health-related quality of life [24–27].

The phenotypes seen in OI exhibit an extensive heterogeneity. The plethora of findings, including oral and craniofacial manifestations, can also differ within the same type of OI.

From birth to adult life, the children and adolescents grow and develop both physically and mentally. All ages are associated with the need for special considerations when it comes to securing oral health and monitoring craniofacial development. A solid base of knowledge is mandatory for diagnosis and treatment of traits. Defining a set of reliable and valid outcomes for individuals with OI that would cover all essential oral and craniofacial aspects and be applicable worldwide is challenging. Such an outcome set could potentially improve the standard of care on both an individual and a population level. Standardization of outcome measures is of utmost importance to enable aggregation of data from different studies, to compare data from different data sets, to allow evidence synthesis, and most importantly, be relevant to the individuals affected by the disorder [28]. Individuals with rare diseases are at increased risk of unmet clinical needs due to limited access to information and clinical care [29, 30]. To meet this challenge, the Care4Brittle-Bones foundation initiated the Key4OI Plus, a project to develop a minimum standard set of outcomes and associated measures for the comprehensive appraisal of OI that would reflect the complexity of OI care and focusses on the issues that are considered most important by individuals with OI [31].

The primary aim of this initiative was to reach an interdisciplinary, international consensus for a standard set of outcomes and measuring instruments for oral healthcare in OI, based on the aspects considered important by both the dental profession and individuals with OI. This standard set is to be comprehensive enough to cover the full range of dental care in OI, as well as practical enough for valid implementation. This approach would permit oral healthcare teams to measure and monitor their performance in a consistent way. Furthermore, this would support longitudinal and cross-sectional comparisons of outcomes between centers that serve OI populations in different countries and cultural contexts.

Methods

Participants of the project team

The Care for Brittle Bones Foundation reached out to all main OI patient organizations, including OIFE and OIF to identify any national experts for oral health/dentistry in their respective countries. In addition, the identified experts reviewed literature to ensure all available expertise was invited to join the project team. The project team was composed of thirteen individuals from eight different

countries, both patient experts and dental care providers with different specialties in dentistry.

The project was led by a project coordinator (n=1) from the Care4BrittleBones Foundation. Clinical disciplines represented included pediatric dentistry (n=5), orthodontics (n=3), oral and maxillofacial surgery (n=1) and prosthodontics (n=1). Furthermore, patient representatives with a professional background in healthcare (n=2) were represented. Of the professional experts, five were involved in dental care for children and five experts involved in dental care for both children and adults.

Initiation of the project

The project was initiated and coordinated by the Care4BrittleBones Foundation. The project used the same approach and principles as for the Key4OI Outcome set [31]. For the original Key4OI work an ethical review was conducted and confirmed that no ethical review was required for the development of outcome measures under this project.

The project commenced with a meeting explaining the purpose and voluntary nature of the project participation. A total of thirteen (monthly and later biweekly) virtual consensus meetings were facilitated by the project coordinator, in the period November 2020–January 2022. All meetings were summarized and recorded, in order to address occasional absences of individual members.

Literature review

In the first phase, the project team collected possible outcome measures related to oral health, dental condition, and occlusion in people with osteogenesis imperfecta, based on available literature and clinical experience. The expert team conducted a non-systematic in-depth literature review of relevant publications in OI and oral health, based on the expertise of each specialist. Variables present in the literature were merged with the extensive collective clinical experience from the expert group. Literature inclusion criteria were original research articles and publications in peer-reviewed journals. Exclusion criteria were articles not available in English or the inability to obtain the full-text article. This resulted in a list of possible clinician-reported outcome measures (CROMs) and expected relevant CROMs. The results were shared in consensus meetings and converted into a comprehensive list of aspects of both CROMs and patient-reported outcome measures (PROMs). Based on the literature, relevant measuring instruments attributing these outcome measures were collected.

Focus groups

The patient representatives and project coordinator recruited individuals with OI to participate in focus

groups. They worked together with various patient organizations to inform the patient community about the upcoming project, and they shared the information on social media. The group consisted of eight male and twelve female patients. The type of OI was distributed as follows: Type I: 6, Type III: 8 and Type IV: 6. To maintain anonymity, complete health records of the focus group participants were not documented. Not all patients consented to record their age. All were above 18 years. The youngest patient was 18 and the oldest 58. The patients were from 11 different countries: two from the US, one from Asia and the others from European origin. The professional experts of the project team were not involved in recruiting the focus groups and did not participate in focus group sessions, to ensure participant anonymity, voluntariness, and unbiasedness.

The project coordinator organized three focus group meetings on 24th of February, 3rd and 9th of March 2021. A total number of 20 adults with OI attended and discussed in the group, supported by a moderator and an innovative IT tool called Mural[®] (see www.mural.com). It is a well-established and proven platform to engage virtual audiences. The meetings were provided by written comments, a summary based on the written comments composed by the two patient experts of the project team, and recordings of the meeting.

During the sessions, 29 themes mentioned by the participants within the sessions, were discussed. Key themes were defined as themes mentioned in all sessions and they were marked as “top priority”. Themes that strongly resonated in two of the three sessions were marked as “priority”.

The key themes mentioned by the focus groups were explained by the patient experts and discussed in project team meetings. The key themes were compared and attributed to the selected CROMs. The items on the CROMs list were discussed, for: relevance of the outcome measure and applicability (time, languages and costs) and validation (also in different languages) of possible measuring instruments. The CROMs mentioned by the focus groups were considered more relevant, than the CROMs mentioned by the experts only.

A different approach to PROMs was adopted. PROMs measuring instruments consisted of mainly questionnaires. All items of the different questionnaires were split into different items/questions. The key themes of the focus groups were matched with the items of the questionnaires, in order to identify the measuring instruments (i.e. questionnaires) covering most and most relevant (high priority) items.

Consensus: Delphi rounds

A modified Delphi technique was used to develop a consensus on a minimal standard outcome set of CROMs and PROMs. The Delphi technique is an iterative multi-stage process to actively transform opinion into group consensus [32]. In a Delphi study, anonymous responses of the project team members are aggregated and shared with the group after each round, ultimately leading to a group consensus. The consensus should be based on the data derived from professional experts and individuals with OI themselves.

In a digital platform, commonly used for scientific surveys, experts anonymously rated the proposed outcome measures and measuring instruments for inclusion in the set, on a 9-point scale. A minimum of 80% with a score of seven, eight or nine was required for the final confirmation of each individual component comprising the measuring instrument. A score of one, two or three in 80% of responses lead to a rejection. Mid-range scores of four, five or six were regarded as “non-conclusive”. The anonymous “non-conclusive” responses were re-discussed in the next consensus meeting and tabled in the next Delphi round. A participation of 80% of experts was required in the Delphi rounds [32, 33].

Results

Literature review and expert opinion/clinical experience

In the first phase of the project, relevant dental items were identified by the project team. The literature review and expert opinion resulted in 35 articles that were reviewed and discussed by the project team. Table 1 outlines the relevant oral health-related aspect and measuring instruments addressing them identified by the professional experts of the project team.

Focus groups

During the sessions, 29 themes were discussed. A total of 13 themes were addressed as relevant to individuals with OI: 6 themes came out as key themes in all sessions (marked as “top priority”) and 7 additional themes (the other dental items) strongly resonated in 2 of the 3 sessions (Table 2). The themes from the focus groups were reviewed and discussed by the professional experts of the project team and were assessed as being covered by the initially identified dental items. The key themes mentioned by the focus groups were attributed to the relevant CROMs and/or listed as a PROM (Table 3).

Table 1 List of possible outcome measures, dental items in osteogenesis imperfecta

Clinician-related outcome measures (CROMS)

- General oral health
- Hard tissues and oral health
 - Periodontal health
 - Tooth wear
 - Plaque
- Osteogenesis imperfecta related oral health
 - Dentinogenesis imperfecta
 - Expressivity—Mild/Moderate/Severe
 - Clinical indicators
 - Radiographic indicators
 - Histological assessment
- Other dental anomalies
 - Shell teeth
 - Tooth agenesis
 - Taurodontism
 - Pulpal stones
 - Ectopic molars (impaction/retention), eruption first molars
 - MIH
 - Other anomalies
- Malocclusion—orthodontics
 - Sagittal incisal relationship
 - Vertical incisal relationship
 - Sagittal molar occlusion
 - Vertical molar occlusion
 - Transversal molar occlusion
 - Occlusion and aesthetics
 - Orthodontic treatment priority
- Osteogenesis imperfecta related (medical) contra-indications or barriers with regard to dental treatment

Patient-reported outcome measures (PROMs)

- Oral pain and dysfunction
- Oral health-related quality of life
- Oral function
 - Aesthetics
- Burden of treatment
- Accessibility to dental treatment
- Anxiety related to dental treatment

Selection of outcome measures and measuring instruments

Based on the input of the professional experts of the project team and the focus groups in the previous phases, outcome measures were formulated, and corresponding measuring instruments were collected (Table 4).

In the five Delphi rounds, carried out between September 2021 and January 2022, a 100% consensus was reached on a set of oral health-related outcome measures and measuring instruments. A clinical practice guideline

Table 2 Key themes identified in focus groups across sections

Key themes identified	Priority
Breaking off tooth pieces, chipping, fracturing teeth	Top
Dental implants	Top
Embarrassed because of teeth or mouth	Top
Ignorance of dentists	Top
Experiencing pain (tooth or jaw)	Top
Proportions of jaw and how they "match" (over/under/open/crossbite)	Top
Affordability of dental treatment	
Anxiety in relation to future of jaws/teeth	
Difficulty eating certain type of food	
Earache	
Facings not sticking/coming off or loosening	
Fillings/Crowns	
Root canal issues (infection, disappearance of visible canal)	

Table 3 Attribution of key themes mentioned by the focus groups to the relevant CROMs and/or listed as a PROM

Key themes identified across sessions	Attribution to CROMs or PROMs
Breaking off tooth pieces, chipping, fracturing teeth	CROMs: OI-related oral health/DGI OI-related oral health or general oral health/tooth wear PROMs: Functional impairments
Dental implants	CROMs: OI-related medical contra-indications or barriers
Embarrassed because of teeth or mouth	CROMs: OI-related oral health/DGI and malocclusion PROMs: Esthetics
Ignorance of dentists	PROMs: separate (new) item
Experiencing pain from teeth or jaws	CROMs: General oral health PROMs: Orofacial pain and dysfunction
Proportions of jaws and how they "match" (over/under/open/crossbite)	CROMs: Malocclusion PROMs: Esthetics Functional impairment?
Affordability of dental treatment	PROMs: separate (new) item
Anxiety in relation to future of jaws / teeth	PROMs: anxiety Discussion: anxiety for dental treatment or insecurity about situation/possibilities/barriers
Difficulty eating certain type of food	CROMs: General oral health (missing teeth, type of prosthetic appliance, periodontal condition) Malocclusion Breaking/chipping > OI-related oral health/DGI PROMs: Orofacial pain and dysfunction
Earache	CROMs and PROMs Discussion: orofacial pain and dysfunction possible aetiology
Facings not sticking / coming off or loosening	CROMs: OI-related oral health / DGI PROMs: Functional impairment
Fillings/Crowns	CROMs: OI-related oral health/DGI
Root canal issues (infection, disappearance of visible canal)	CROMs: OI-related oral health/radiographic DGI indicators/pulp obliteration

Table 4 Selection of outcome measures, with input of the focus groups and possible measuring instruments

Domain		Outcome measure	Measuring instruments	
<i>CROMs</i>				
General oral health		Hard tissues and oral health	DMFT (Decayed Missing Filled Teeth) [65]	
		Periodontal health	BPE (Basic Periodontal Examination) CPITN (Community Periodontal Index of Treatment Needs) Periodontal status [34–36, 66, 67]	
		Endodontic health (periapical pathology)	Clinical and/or radiographic assessment	
		Tooth wear	BEWE (Basic Erosive Wear Examination) [68] TWI (tooth wear index) [69]	
		Plaque	Plaque score [37]	
		Number of permanent teeth	Clinical or radiographic assessment	
		Number of extracted teeth	Clinical or radiographic assessment	
		Number of teeth with direct/indirect restorations	Clinical and/or radiographic assessment	
		Number of dental implants	Radiographic assessment	
		Number of endodontically treated teeth	Radiographic assessment	
		Type of dental prosthesis	Clinical assessment: 0. Implant supported crowns 1. Fixed partial denture, dental supported 2. Fixed partial denture, implant supported 3. Removable partial denture 4. Full denture 5. Other?	
	OI specific oral health /DGI spectrum	DGI—Expressivity	Expressivity	Clinical and radiographic assessment: 3 types of expressivity: 1. Presence in primary dentition 2. Presence in both 3. Isolated histological DGI only
		DGI—Clinical DGI indicators	1. Pathologic discoloration	Clinical assessment—according to scale or description (blue/grey or yellow/brown) [8, 14, 38]
2. Attrition			BEWE [68] TWI [69]	
3. Fractures			Clinical assessment	
DGI—Radiographic DGI indicators		1. Bulbous crowns with cervical constriction	Radiographic assessment (Y/N) [8]	
		2. Pulpal obliteration	Radiographic assessment (Y/N AND partial (Pulp chamber is not visible and the canal is markedly narrowed but visible) or total (the pulp chamber and canal is hardly or not visible))	
DGI—Histological assessment		3. Small radices / short roots Histological DGI	Radiographic assessment Histological examination [60]	

Table 4 (continued)

Domain	Outcome measure	Measuring instruments	
OI specific oral health	Shell teeth	Clinical and radiographic assesment	
	Tooth agenesis	TAC (Tooth Agenesis Code) [70] Hypodontia/oligodontia Registration of the number and specification of teeth congenitally absent [22, 71]	
	Taurodontism	Radiographic assessment (first molars) (Y/N) [12, 69]	
	Pulpal stones	Radiographic assessment (Y/N)	
	Permanent molar eruption (impaction or retention)	Relevant if it occurs in association with DGI and/or aberrant craniofacial development/malocclusion (Y/N in association with DGI and/or aberrant craniofacial development/malocclusion) Clinical and radiographic assessment. Categorizing by categories: 1. Developing tooth 2. Normal eruption 3. Mesioangular impaction of maxillary molar 4. Retention of maxillary molar (not erupting, no physical obstacles eruption pathway) 5. Impaction of mandibular molar (due to ectopic position or an obstacle) 6. Retention of mandibular molar (not erupting, no physical obstacles in eruption pathway) [72, 73]	
	Molar-incisor-hypomineralisation	EAPD classification [74]	
	Other anomalies	Clinical and/or radiographic assessment	
	Craniofacial / Orthodontics	Sagittal incisal relationship	Clinical assessment (overjet in mm OR maxillary overjet (> 5 mm) Y/N and mandibular overjet (=0 mm or <0 mm) (Y/N) [75]
		Vertical incisal relationship	Clinical assessment (assesment of overbite in mm OR anterior open bite (< 0 mm) Y/N and deep bite (> 4 mm) Y/N)
		Sagittal molar occlusion	Clinical assessment (occlusion according to Angle classification)
Vertical molar occlusion		Clinical assessment (lateral open bite in molar region Y/N)	
Vertical molar occlusion		Clinical assessment (lateral open bite in premolar region Y/N)	
Transversal molar occlusion		Clinical assessment (crossbite Y/N? Crossbite uni- or bilateral)	
Orthodontic diagnosis		Radiographic assessment? (ortho-pantomography, cephalometry, Cone beam CT)	
Occlusion and aesthetics		ICON (Index of Complexity, Outcome and Need) [76]	
Orthodontic treatment priority	IOTN (index on treatment need and indication of why the treatment need existst) [77]		

Table 4 (continued)

Domain	Outcome measure	Measuring instruments
<i>PROMs</i>		
Pain and dysfunction	Pain, temporomandibular disorders, orofacial pain and dysfunction	DC/TMD [78, 79] 3 questions TMD screening [48, 49] MFIQ [80]
OHRQoL/ aesthetics	Oral function	COHIP-SF 19 [81] CPQ [46, 47, 82] OHIP-49 [45] OHIP-14 [83] according to PROMS AI (SP) [84]
Burden of (dental) treatment	Aesthetics	OES (Orofacial Esthetic Scale) [85, 86] No. of appointments, duration, patient perception
Dental anxiety	Anxiety (related to dental treatment)	CEDAM [87] MCDAS [88, 89] Single question with VAS [90]
Accessibility to dental treatment (input focus groups)		–

recommendation that was considered not to be part of the set but too important to be neglected, was also made. This recommendation was included based on acceptance for inclusion by 10 out of 12 experts (> 80%). The recommended set of outcome measures and measuring instruments is represented in Table 5.

Clinician-related outcome measures (CROMS)

The project team recommended to measure general oral health and OI-specific oral health items.

General oral health

General oral health is recommended to be measured as the number of decayed, missing and filled teeth (DMFT for permanent dentition and dmft for deciduous teeth) and the periodontal health with the screening tools of Basic Periodontal Examination (BPE) and Community Periodontal Index (CPI) [34–36]. In addition, plaque score was included in the assessment of general oral health [37]. Recommended frequency for recording of DMFT was in conjoint with every routine examination but at least every second year, and dmft in the primary or mixed dentition at least at age 3, 6 and 13 years. In addition, consensus was reached for the need to record reasons for the “filled” as either: (1) Tooth wear and/or chipping and/or fracture (non-cariogenic tooth substance loss), (2) Caries and (3) Unknown aetiology. The reasons behind the DMFT/dmft numbers were considered more important and indicative for specific oral problems, than the absolute number.

Measuring periodontal health by Basic Periodontal Evaluation (BPE) from age 7 at every routine examination, was recommended.

OI-specific oral health

Measurement of OI-specific oral health was divided into 5 items: DGI, other (dental) anomalies, items related to occlusion, tooth wear, and OI-related medical contraindications or barriers regarding dental treatment.

Dentinogenesis imperfecta

The presence (yes/no) of DGI should be recorded, based on clinical and/or radiographic indicators [38]. As the presence of DGI does not change with time, but the expressivity may differ significantly between the primary and permanent dentition, two baseline assessments were recommended: first for the primary dentition between 3 and 6 years of age at latest, and second in the permanent dentition, when all the permanent teeth except for wisdom teeth, are present [12].

Other dental anomalies

Other dental anomalies or relevant dental findings should be recorded. Special emphasis should be put on presence of tooth agenesis, shell teeth (as an early sign of DGI before obliteration of the pulp), taurodontism, ectopic permanent second molars (impaction/retention) and ectopic eruption of first molars, as these anomalies are more prevalent in individuals with OI compared to the general population [9, 11, 12, 22, 39].

Table 5 Standard set of outcome measures and measuring instruments for the assessment of oral health and occlusion of individuals with Osteogenesis Imperfecta

General oral health	Hard tissues and oral health	<p>Adults: DMFT* every routine examination, at least every 2nd year Children: dmft* at least at age 3, 6, 13 (DMFT when permanent teeth have erupted) *Including: reasons for "filled" as follows: (1) tooth wear and/or chipping and/or fracture (non-cariogenic tooth substance loss) (2) caries (3) unknown aetiology</p>
OI-specific oral health	<p>Periodontal health Dentinogenesis imperfecta Dental anomalies Occlusion</p>	<p>BPE, from age 7 at every routine examination Y/N (clinical/radiological), 2 baselines: primary (3–6 years of age latest) and permanent (when all have erupted) Recording of anomalies (shell teeth, tooth agenesis, taurodontism, ectopic permanent second molars (impaction/retention), eruption of first molars) Horizontal overjet Mandibular overjet Vertical overbite Anterior open bite</p> <p>Horizontal overjet: measurement (if overjet ≥ 0 mm) and clinical assessment of presence of increased horizontal overjet (overjet > 5 mm) Y/N Mandibular overjet: clinical assessment of presence of mandibular overjet (overjet ≤ 0 mm Y/N) and measurement (if mandibular overjet ≤ 0 mm) Vertical overbite: measurement of vertical overbite (if VOB ≥ 0 mm) and clinical assessment of presence of increased vertical overbite (≥ 5 mm) / deep bite Y/N Anterior open bite: clinical assessment (VOB < 0 mm Y/N) (3) and measurement (if VOB < 0 mm)</p>
	Crossbite Posterior open bite	<p>Crossbite: clinical assessment of presence of crossbite (Y/N) and notification of crossbite type 1 (Unilateral / bilateral / anterior) and/or crossbite type 2 (Molar/premolar-canine / incisor) Posterior open bite: clinical assessment of presence of crossbite (Y/N) and notification of posterior open bite type 1 (Unilateral / bilateral) and/or posterior open bite type 2 (Molar/premolar-canine)</p>
	Clinical assessment of molar eruption Any kind of malocclusion	<p>Clinical assessment of molar eruption Any kind of malocclusion should be recorded. Special consideration should be put on presence of the previously mentioned variables as they are more prevalent in individuals with OI</p>
	Tooth wear	<p>Identification of the presence of pathological tooth wear (i.e. non-physiological) by recording " (1) yes, (2) possibly, (3) no/physiological When screening for pathological tooth wear make use of the appropriate methods suitable for the specific situation in an individual patient. The use of indices, photographs, study casts, digital 3Ddatasets, clinical examination and anamnestic information can be part of that</p>
OI-related (medical) contra-indications or barriers with regard to dental Tx	Recording medical treatment with bisphosphonates If bisphosphonates are used: record start and end date of treatment Prior dental procedures/treatment: Clinical assessment and history taking	

Table 5 (continued)

Oral Health Related Quality of Life (OHRqOL)	We recommend measurement of Oral Health Related Quality of Life by CPQ 8–10 at age 8 and CPQ11–14 at age 11
Temporomandibular Disorder (TMD)	We recommend measurement of Oral Health Related Quality of Life by OHRP-49, starting at age 15, every 5 years in children and adolescents, we recommend measurement (screening) of TMD-problems by 3Q/TMD annually, starting at age 10 In adults (above 19 years), we recommend measurement (screening) of TMD-problems by 3Q/TMD with an interval of 2 years
Basilar invagination	If radiographic scans (CBCT or Ceph) are made or already exist, the dentist has a responsibility that any anomalies or pathologies are reviewed, including and specifically concentrating on any indications for BI. It is recommended to refer to a radiologist with relevant competence for this assessment

Occlusion

Malocclusion is a common finding in individuals with OI. With regard to occlusion, measurement of horizontal overjet and vertical overbite is recommended, as well as recording the presence of possible open bite, crossbite, ectopic eruption of teeth and/or other occlusal deviation [40, 41].

Tooth wear

Consensus was reached on the relevance of the assessment of tooth wear in individuals with OI. Little evidence is available, but tooth wear in combination with DGI with accompanying risk for chipping and fractures and the compromised condition of dentin, was presumed to be an important factor in tooth prognosis. No consensus was reached on the index that should be used. However, recommendation included identification of the presence of pathological (i.e. non-physiological) tooth wear by recording: (1) Yes; (2) Possibly; (3) No/physiological. As no consensus could be reached, it was recommended to make use of the appropriate methods suitable for the specific situation in the individual patient. The use of indices, photographs, study casts, digital 3D datasets, clinical examination and anamnestic information can be part of that.

OI-related medical contra-indications or barriers with regard to dental treatment

Prior to dental procedures or treatment, clinical assessment and history taking should be performed. Medical contra-indications or barriers should be recorded. Especially medical treatment with bisphosphonates is considered relevant. If bisphosphonates are used, start and end date should be recorded. The reason is the potential effects of bisphosphonates on tooth movement in orthodontic treatment, tooth development and medication-related osteonecrosis of the jaw (MRONJ) [42–44].

Patient-reported outcome measures (PROMs)

It was recommended to measure oral health-related quality of life by CPQ8-10 at age 8, CPQ11-14 at age 11 and OHIP-49 starting at age 15, every 5 years [45–47].

Consensus was reached on measuring (screening) temporomandibular disorders (TMD), by 3 screening questions (3Q/TMD) [48–50]. Screening was recommended to be performed annually starting age 10, and above 19 years with an interval of 2 years.

The Delphi results on ‘anxiety’ were inconclusive. Discussion on this topic led to no views on this topic and anxiety was not included in the final set.

Recommendation

The importance of assessment of basilar invagination or impression was acknowledged in several discussion meetings. Basilar invagination is a serious co-morbidity of OI that may cause life-threatening compression of medulla and cervical spine [51–53]. Lateral skull radiographs, Conebeam CT (CBCT), or MR/CT-images used for dental/orthodontic assessment (cephalometrics) can reveal asymptomatic cranial base pathologies [52, 54]. However, no consensus could be reached on inclusion and especially measuring methods of BI in this set of oral health-related outcome measures. In the final Delphi, consensus was reached on the following recommendation: If radiographic scans (CBCT or Cephalometric radiographs) are obtained or already exist in patient records, the dentist has a responsibility that in case any anomalies or pathologies are observed, to refer the patient to a radiologist with relevant competence for assessment of craniocervical pathology.

Discussion

In this Delphi consensus study, we developed a standard set of outcome measures and measuring instruments on oral health and occlusion in individuals with OI, which can be implemented by healthcare professionals all over the world. The standard set of outcome measures proposed in this study enables the assessment and comparison of relevant dental and oral health problems. In addition, systematic implementation of a standard set of outcome measures by oral healthcare professionals can facilitate future research on dental and oral health problems in people with OI.

At present, several centers providing care for individuals with OI have established consensus guidelines on the use of bisphosphonate therapy, physical rehabilitation and surgical management of fractures [55–57]. This is the first oral health-focused guideline project that incorporates patient-reported outcome measures into clinician-related outcome measures reviewed by a panel of individuals with OI and an international group of experts.

Although not included in the Delphi rounds, it is important to note that an oral hygiene inventory should be conducted for all patients during each visit. A consistent daily oral hygiene routine has a substantial impact on individual’s oral health and affects the other outcome variables. The frequency of routine examination was not included in the standard set. The authors acknowledge that regular follow-up on an individually based frequency is of importance, especially when factors are present that may hamper oral health such as a more severe phenotype.

In people with OI, especially when multiple health issues are present, assessment of the oral health may not obtain highest priority [58]. However, the outcome of

the focus groups sessions emphasized the importance of dental and oral problems as well as the impact they have on everyday life, especially when oral functional impairments or esthetic problems are present. Focus group discussions stated that dental problems and oral health-related concerns can be left underestimated and that oral health professionals may appear reluctant to act on them. The absence of universal treatment guidelines, limited scientific evidence and knowledge, and limited clinical experience in a rare disorder like OI may contribute to this reluctance [59].

Previous studies have demonstrated that the dental concerns of children and adolescents with OI affect functional and socio-emotional well-being and thereby oral health-related quality of life [24, 27]. This emphasizes the relevance of adequate attention for oral health and oral health-related well-being, in individuals with OI.

Strengths and limitations

A strength of this study is that the consensus statements were based on input from both focus groups and oral healthcare providers. The input from the focus groups formed the basis for the set of outcome measures. Moreover, the selection of the specific measuring instrument for assessment of oral health-related quality of life, was derived from the focus groups input. The oral healthcare providers, all have a special interest in OI and covered different areas of specialties of dentistry relevant for full assessment in complex dental/oral conditions. The participation of two patient experts, both professionals in healthcare, was considered very valuable. The geographic representation of the experts can be considered a limitation: the expert team involved professionals from Europe, North America and Asia but none from Africa, Oceania, or South America. Another limitation of the study is that a certain degree of computer skills was required to attend the focus group sessions possibly affecting the group composition. Illiterate people and children were not represented. It is possible that those individuals with OI, who experience oral health-related problems were more prone to participate in the focus groups. This possible sampling bias can be also considered a strength of the study as it could be speculated to result in addressing of the relevant dental problems. However, it might also imply a biased view on the dental problems when people without dental problems or positive experience did not participate.

Two of the priority items mentioned in the focus group sessions, were not selected as an outcome measure by the oral health professionals. Both *ignorance of dentists* and *affordability of dental treatment* were rejected based on discussion in the expert team sessions, due to lack of universal methods to measure them for global comparison.

Nevertheless, these items are highly important to people with OI.

To date, no OI specific questionnaire exists for Oral Health. Hence, the assessments included in the standard set are not validated for OI. OI is a very heterogeneous disorder and the prevalence is low. Therefore, to develop and implement a validated OI specific questionnaire for Oral Health in OI would be very challenging. The fact that the disorder is unknown for most oral health professionals emphasizes the importance of our study in raising awareness and understanding of oral health issues associated with OI. This study aimed to do this initially using validated non-OI specific questionnaires, as this approach is scientifically more sound in the short and mid-term. In the long term, developing an OI-specific validated questionnaire is certainly a topic for future research. DGI may be diagnosed by clinical, radiographic and/or histologic findings. In cases of no obvious clinical or radiographic findings, a histologic examination of an exfoliated or therapeutically extracted tooth may still reveal dentin anomalies associated with DGI [12, 60]. Based on this extensive phenotypic heterogeneity (clinically, radiographically, histologically), the expert group discussion led to the suggestion for a more comprehensive classification of OI-related DGI. This would facilitate diagnosis and increase the basis for well-founded treatment guidelines. However, the opportunity for histological examination differs between countries. Based on this, the expert group considered it as not being part of the minimum standard outcome set. The need for a classification of OI-related DGI was acknowledged but considered beyond the scope of this study.

At present, no OI-specific tooth wear index exists that would take into consideration the presence and effect of DGI. In individuals with OI and DGI, the enamel frequently chips from the affected dentin leading to tooth fractures [61]. There is no gold standard in assessment of tooth wear. Although recommendations exist on identification and treatment of pathological tooth wear, it might not be suitable for use in people with DGI [62, 63].

The recommended frequency for recording of DMFT was in conjoint with every routine examination but at least every second year, and dmft in the primary or mixed dentition at least at age 3, 6, and 13 years. In case of adults without registrations and when the cause of missing teeth is not remembered, the adjusted decayed and filled teeth (ADFT) can be applied, to overcome this problem [59, 64].

In the final set of outcome measures and measuring instruments both measurement of horizontal overjet and mandibular overjet are included. Mandibular overjet can be interpreted and measured as a negative horizontal overjet (frequently seen in OI) as well. The same accounts

for vertical overbite and anterior open bite: the latter can be measured as a negative vertical overbite.

The validated OHIP-49 questionnaire on oral health related quality of life was included in our standard set, instead of the more compact OHIP-14. This choice was based on the suggestion of the patient experts in our project team, with argumentation that the questions in this more comprehensive version reflected more of the issues that people with OI encounter. Both instruments are validated on the oral health related quality of life. If a more efficient tool for a quick assessment is required, OHIP-14 may be sufficient.

The use of a validated questionnaire to evaluate treatment will provide essential information on the impact of specific interventions. However, this was not included in this standard set.

'Pain' was mentioned as a top-priority item, during the focus group sessions. Participants mentioned tooth and jaw pain, but also earache. Pain is complex and can have multiple causes and other factors may contribute to, amplify or interfere with pain sensation. Without further specification and assessment of the pain, it was difficult to address it. In dentistry most common origins of pain are odontogenic pain (dental pain, pain of the teeth or surrounding tissues) or pain related to temporomandibular dysfunction.

A limitation of this study is the lack of children focus groups. Focus groups can successfully include children. However, it also creates challenges with e.g. need of adaptations based on maturity and age of the included children. Moreover, ensuring a comfortable and safe environment for children to express their opinions despite the power imbalances between adults and participants is of utmost importance. These issues need special competence of the investigating team with care taken to required considerations. The team stresses the importance of future studies investigating variables of importance for children's oral health related quality of life. Children focus groups would then be of significant value.

The authors acknowledge that in this study not all continents and countries were represented. The project initiators reached out to any known expert who had published about the topic, any expert center worldwide and any expert known to the patient organizations. Many experts have very small populations and their interest was often not significant enough to join the project or despite interest there were competing priorities as OI was only one out of many disorders they supported. The authors recognize that it is most essential that dentists are aware of OI and the consequences for oral health. It should also be underlined that there is a need of continuing networking to expand

the knowledge of available expert teams and to enhance the collaborations between countries in the oral health care in OI, but also in other rare diseases.

Conclusion

In this Delphi consensus study, a standard set of outcome measures and measuring instruments was developed for identification of prioritized oral health-related problems in OI. It is recommended to be implemented by dental practitioners in order to standardize and equalize the dental care in children and adults with OI. The minimum set of outcome measures ensures feasibility of use and requires only limited time from the clinician.

The use of a standardized set of outcome measures and measuring instruments will also facilitate future research and collaboration.

This is a first important step in composing a standard for OI specific oral healthcare. Further research and collaboration are necessary to identify and specify the oro-dental symptoms and problems of people with OI in broader perspective and eventually compose treatment guidelines, in order to improve the quality of care and oral health-related quality of life in individuals with osteogenesis imperfecta.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13023-024-03308-5>.

Supplementary Material 1

Author contributions

LB, HA, AA, SC, HG, RJ, ML, DM, SP, JMR, TS, LZ, KA. The project was initiated by DM. DM organized the online meetings of the Delphi study, led the three focus group sessions, facilitated and hosted the Delphi rounds, analysed the results of the Delphi rounds. TS and SC participated in the focus groups. SC summarized the outcomes of these sessions with consent of participants of the focus groups. Authors LB, HA, AA, HG, RJ, ML, SP, JMR, LZ and KA collected and read articles, related to oral health and occlusion relevant for osteogenesis imperfecta and derived possible outcome measures and measuring instruments from this. Authors LB, HA, AA, SC, HG, RJ, ML, SP, JMR, TS, LZ and KA participated in the Delphi rounds. LB was penholder during the process. LB and KA interpreted the results of the Delphi rounds and LB, KA and DM formulated the statements of the following Delphi round. LB, HA and KA drafted the manuscript. All authors, LB, HA, AA, SC, HG, RJ, ML, DM, SP, JMR, TS, LZ and KA read and approved the final manuscript.

Funding

The project was done for the most part as a volunteer effort of all parties involved. Limited funding was provided by the not-for-profit Care4BrittleBones foundation based on crowdfunding in the OI-community (no funding from industry). The funder had no involvement in analysis and interpretation of data.

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Declarations

Ethics approval and consent to participate

For the original Key4OI work an ethical review was conducted and confirmed that no ethical review was required for the development of outcome measures under this project.

Consent for publication

All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Vogellanden, Center of Rehabilitation Medicine and Special Care in Dentistry, Zwolle, The Netherlands. ²Department of Oral and Maxillofacial Diseases, University of Helsinki, Helsinki, Finland. ³Helsinki University Hospital Head and Neck Center, Helsinki, Finland. ⁴Whittington Health NHS Trust Dental Services, London, UK. ⁵Care4BrittleBones Foundation, Wassenaar, The Netherlands. ⁶Elkerliek Hospital, Helmond, The Netherlands. ⁷Center for Oral Health in Rare Diseases, Department of Dental and Maxillofacial Surgery, Aarhus University Hospital, Aarhus, Denmark. ⁸Department of Paediatric Dentistry, University of Bristol, Bristol, UK. ⁹Department of Stomatology, The University of Hong Kong-Shenzhen Hospital, Shenzhen, China. ¹⁰Department of Paediatric Dentistry, UCL Eastman Dental Institute, London, UK. ¹¹Department of Molecular Genetics, Baylor School of Medicine, Houston, TX, USA. ¹²ProRaris, Vuarrens, Switzerland. ¹³Mun-H-Center, National Orofacial Resource Centre for Rare Diseases and Clinic of Pedodontics, Public Dental Service, Region Västra Götaland, Göteborg, Sweden. ¹⁴Department of Pediatric Dentistry, Institute of Odontology, Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden. ¹⁵Department of Dental Medicine, Division of Orthodontics and Pediatric Dentistry, Karolinska Institutet and Center for Pediatric Oral Health Research, Stockholm, Sweden.

Received: 18 October 2023 Accepted: 6 August 2024

Published online: 13 August 2024

References

1. Marini JC, Forlino A, Bächinger HP, Bishop NJ, Byers PH, Paepe A, Fassier F, Fratzl-Zelman N, Kozloff KM, Krakow D, Montpetit K, Semler O. Osteogenesis imperfecta. *Nat Rev Dis Primers*. 2017;18(3):17052.
2. Sillence DO, Rimoin DL, Danks DM. Clinical variability in osteogenesis imperfecta-variable expressivity or genetic heterogeneity. *Birth Defects Orig Artic Ser*. 1979;15(5B):113–29.
3. Sykes B, Ogilvie D, Wordsworth P, Anderson J, Jones N. Osteogenesis imperfecta is linked to both type I collagen structural genes. *Lancet*. 1986;2(8498):69–72.
4. Wallis GA, Sykes B, Byers PH, Mathew CG, Viljoen D, Beighton P. Osteogenesis imperfecta type III: mutations in the type I collagen structural genes, COL1A1 and COL1A2, are not necessarily responsible. *J Med Genet*. 1993;30(6):492–6.
5. Marini JC, Forlino A, Cabral WA, Barnes AM, San Antonio JD, Milgrom S, Hyland JC, Körkkö J, Prockop DJ, De Paepe A, Coucke P, Symoens S, Glorieux FH, Roughley PJ, Lund AM, Kuurila-Svahn K, Hartikka H, Cohn DH, Krakow D, Mottes M, Schwarze U, Chen D, Yang K, Kuslich C, Troendle J, Dalgleish R, Byers PH. Consortium for osteogenesis imperfecta mutations in the helical domain of type I collagen: regions rich in lethal mutations align with collagen binding sites for integrins and proteoglycans. *Hum Mutat*. 2007;28(3):209–21.
6. Van Dijk FS, Sillence DO. Osteogenesis imperfecta: clinical diagnosis, nomenclature and severity assessment. *Am J Med Genet A*. 2014;164A(6):1470–81. Epub 2014 Apr 8. Erratum in: *Am J Med Genet A*. 2015;167A(5):1178.
7. Unger S, Ferreira CR, Mortier GR, Ali H, Bertola DR, Calder A, Cohn DH, Cormier-Daire V, Girisha KM, Hall C, Krakow D, Makitie O, Mundlos S, Nishimura G, Robertson SP, Savarirayan R, Sillence D, Simon T, Sutton VR, Warman ML, Superti-Furga A. Nosology of genetic skeletal

- disorders: 2023 revision. *Am J Med Genet A*. 2023;191(5):1164–209. <https://doi.org/10.1002/ajmg.a.63132>. (Epub 2023 Feb 13).
8. O’Connell AC, Marini JC. Evaluation of oral problems in an osteogenesis imperfecta population. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1999;87(2):189–96.
9. Malmgren B, Norgren S. Dental aberrations in children and adolescents with osteogenesis imperfecta. *Acta Odontol Scand*. 2002;60(2):65–71.
10. Saeves R, Lande Wekre L, Ambjørnsen E, Axelsson S, Nordgarden H, Storhaug K. Oral findings in adults with osteogenesis imperfecta. *Spec Care Dentist*. 2009;29(2):102–8.
11. Thuesen KJ, Gjørup H, Hald JD, Schmidt M, Harsløf T, Langdahl B, Haubek D. The dental perspective on osteogenesis imperfecta in a Danish adult population. *BMC Oral Health*. 2018;18(1):175.
12. Andersson K, Dahllöf G, Lindahl K, Kindmark A, Grigelioniene G, Åström E, Malmgren B. Mutations in COL1A1 and COL1A2 and dental aberrations in children and adolescents with osteogenesis imperfecta: a retrospective cohort study. *PLoS ONE*. 2017;12(5): e0176466.
13. Lukinmaa PL, Ranta H, Ranta K, Kaitila I. Dental findings in osteogenesis imperfecta: I. Occurrence and expression of type I dentinogenesis imperfecta. *J Craniofac Genet Dev Biol*. 1987;7(2):115–25.
14. de La Dure-Molla M, Philippe Fournier B, Berdal A. Isolated dentinogenesis imperfecta and dentin dysplasia: revision of the classification. *Eur J Hum Genet*. 2015;23(4):445–51 (Epub 2014 Aug 13).
15. Kim JW, Hu JC, Lee JI, Moon SK, Kim YJ, Jang KT, Lee SH, Kim CC, Hahn SH, Simmer JP. Mutational hot spot in the DSPP gene causing dentinogenesis imperfecta type II. *Hum Genet*. 2005;116(3):186–91. <https://doi.org/10.1007/s00439-004-1223-6>. (Epub 2004 Dec 8).
16. Beattie ML, Kim JW, Gong SG, Murdoch-Kinch CA, Simmer JP, Hu JC. Phenotypic variation in dentinogenesis imperfecta/dentin dysplasia linked to 4q21. *J Dent Res*. 2006;85(4):329–33.
17. Barron MJ, McDonnell ST, Mackie I, Dixon MJ. Hereditary dentine disorders: dentinogenesis imperfecta and dentine dysplasia. *Orphanet J Rare Dis*. 2008;20(3):31.
18. Andersson K, Malmgren B, Åström E, Dahllöf G. Dentinogenesis imperfecta type II in Swedish children and adolescents. *Orphanet J Rare Dis*. 2018;13(1):145.
19. Lukinmaa PL, Ranta H, Ranta K, Kaitila I, Hietanen J. Dental findings in osteogenesis imperfecta: II. Dysplastic and other developmental defects. *J Craniofac Genet Dev Biol*. 1987;7(2):127–35.
20. Jensen BL, Lund AM. Osteogenesis imperfecta: clinical, cephalometric, and biochemical investigations of OI types I, III, and IV. *J Craniofac Genet Dev Biol*. 1997;17(3):121–32.
21. Waltimo-Sirén J, Kolkka M, Pynnönen S, Kuurila K, Kaitila I, Kovero O. Craniofacial features in osteogenesis imperfecta: a cephalometric study. *Am J Med Genet A*. 2005;133A(2):142–50.
22. Malmgren B, Andersson K, Lindahl K, Kindmark A, Grigelioniene G, Zachariadis V, Dahllöf G, Åström E. Tooth agenesis in osteogenesis imperfecta related to mutations in the collagen type I genes. *Oral Dis*. 2017;23(1):42–9. <https://doi.org/10.1111/odi.12568>. (Epub 2016 Sep 13).
23. Gjørup H, Jacobsen PE, Hald JD, Haubek D. Craniofacial morphology in adults with osteogenesis imperfecta: a cross-sectional study. *Orthod Craniofac Res*. 2023;26(2):248–55 (Epub 2022 Sep 8).
24. Najirad M, Ma MS, Rauch F, Sutton VR, Lee B, Retrouvey JM; Members of the BBD; Esfandiari S. Oral health-related quality of life in children and adolescents with osteogenesis imperfecta: cross-sectional study. *Orphanet J Rare Dis*. 2018;13(1):187.
25. Najirad M, Madathil SA, Rauch F, Sutton VR, Lee B, Retrouvey JM; Members of the Brittle Bone Diseases Consortium; Esfandiari S. Malocclusion traits and oral health-related quality of life in children with osteogenesis imperfecta: A cross-sectional study. *J Am Dent Assoc*. 2020;151(7):480–490.
26. Gjørup H, Beck-Nielsen SS, Hald JD, Haubek D. Oral health-related quality of life in X-linked hypophosphataemia and osteogenesis imperfecta. *J Oral Rehabil*. 2021;48(2):160–8 (Epub 2020 Nov 19).
27. Cachia Mintoff JM, Riddington M, Parekh S. Oral health-related quality of life in children with osteogenesis imperfecta. *Eur Arch Paediatr Dent*. 2022;23(2):261–70.
28. Williamson PR, Altman DG, Blazeby JM, Clarke M, Devane D, Gargon E, Tugwell P. Developing core outcome sets for clinical trials: issues to consider. *Trials*. 2012;6(13):132.

29. Rousseau M, Retrouvey JM, Members of the Brittle Bone Disease Consortium. Osteogenesis imperfecta: potential therapeutic approaches. *PeerJ*. 2018;17(6): e5464.
30. Slade A, Isa F, Kyte D, Pankhurst T, Kerecuk L, Ferguson J, Lipkin G, Calvert M. Patient reported outcome measures in rare diseases: a narrative review. *Orphanet J Rare Dis*. 2018;13(1):61.
31. Nijhuis W, Franken A, Ayers K, Damas C, Folkestad L, Forlino A, Fraschini P, Hill C, Janus G, Kruse R, Lande Wekre L, Michiels L, Montpetit K, Panzeri L, Porquet-Bordes V, Rauch F, Sakkers R, Salles JP, Semler O, Sun J, To M, Tosi L, Yao Y, Yeung EHK, Zhytnik L, Zillikens MC, Verhoef M. A standard set of outcome measures for the comprehensive assessment of osteogenesis imperfecta. *Orphanet J Rare Dis*. 2021;16(1):140.
32. Boulkedid R, Abdoul H, Loustau M, Sibony O, Alberti C. Using and reporting the Delphi method for selecting healthcare quality indicators: a systematic review. *PLoS ONE*. 2011;6(6):e20476 (**Epub 2011 Jun 9**).
33. Biggane AM, Williamson PR, Ravaud P, Young B. Participating in core outcome set development via Delphi surveys: qualitative interviews provide pointers to inform guidance. *BMJ Open*. 2019;9(11): e032338.
34. WHO. Oral health surveys: basic methods. 5th ed. Geneva: World Health Organization; 2014.
35. BSP: The Basic Periodontal Examination (BPE). 2011 The British Society of Periodontology. Available at: <http://www.bsperio.org.uk/publications/>
36. Cole E, Ray-Chaudhuri A, Vaidyanathan M, Johnson J, Sood S. Simplified basic periodontal examination (BPE) in children and adolescents: a guide for general dental practitioners. *Dent Update*. 2014;41(4):328–30.
37. O’Leary TJ, Drake RB, Naylor JE. The plaque control record. *J Periodontol*. 1972;43(1):38.
38. Shields ED, Bixler D, El-Kafrawy AM. A proposed classification for heritable human dentine defects with a description of a new entity. *Arch Oral Biol*. 1973;18(4):543–53.
39. Andersson K, Malmgren B, Åström E, Nordgren A, Taylan F, Dahllöf G. Mutations in COL1A1/A2 and CREB3L1 are associated with oligodontia in osteogenesis imperfecta. *Orphanet J Rare Dis*. 2020;15(1):80.
40. Rizkallah J, Schwartz S, Rauch F, Glorieux F, Vu DD, Muller K, Retrouvey JM. Evaluation of the severity of malocclusions in children affected by osteogenesis imperfecta with the peer assessment rating and discrepancy indexes. *Am J Orthod Dentofac Orthop*. 2013;143(3):336–41.
41. Retrouvey JM, Taqi D, Tamimi F, Dagdeviren D, Glorieux FH, Lee B, Hazboun R, Krakow D, Sutton VR, Members of the BBD Consortium. Oro-dental and cranio-facial characteristics of osteogenesis imperfecta type V. *Eur J Med Genet*. 2019;62(12):103606 (**Epub 2018 Dec 26**).
42. Friedrich RE, Scheuer HA, Hölzje W. The effect of bisphosphonate medication on orthodontics and orthognathic surgery in patients with osteogenesis imperfecta. *GMS Interdiscip Plast Reconstr Surg DGPW*. 2019;8:Doc06.
43. Contaldo M, Luzzi V, Ierardo G, Raimondo E, Boccellino M, Ferati K, Bexheti-Ferati A, Inchingolo F, Di Domenico M, Serpico R, Polimeni A, Bossù M. Bisphosphonate-related osteonecrosis of the jaws and dental surgery procedures in children and young people with osteogenesis imperfecta: a systematic review. *J Stomatol Oral Maxillofac Surg*. 2020;121(5):556–62 (**Epub 2020 Mar 7**).
44. Malmgren B, Thesleff I, Dahllöf G, Åström E, Tsilingaridis G. Abnormalities in tooth formation after early bisphosphonate treatment in children with osteogenesis imperfecta. *Calcif Tissue Int*. 2021;109(2):121–31 (**Epub 2021 Mar 20**).
45. Slade GD, Spencer AJ. Development and evaluation of the Oral Health Impact Profile. *Community Dent Health*. 1994;11(1):3–11.
46. Jokovic A, Locker D, Stephens M, Kenny D, Tompson B, Guyatt G. Validity and reliability of a questionnaire for measuring child oral-health-related quality of life. *J Dent Res*. 2002;81(7):459–63.
47. Jokovic A, Locker D, Tompson B, Guyatt G. Questionnaire for measuring oral health-related quality of life in eight- to ten-year-old children. *Pediatr Dent*. 2004;26(6):512–8.
48. Lövgren A, Häggman-Henrikson B, Visscher CM, Lobbezoo F, Marklund S, Wänman A. Temporomandibular pain and jaw dysfunction at different ages covering the lifespan: a population based study. *Eur J Pain*. 2016;20(4):532–40 (**Epub 2015 Aug 27**).
49. Lövgren A, Visscher CM, Häggman-Henrikson B, Lobbezoo F, Marklund S, Wänman A. Validity of three screening questions (3Q/TMD) in relation to the DC/TMD. *J Oral Rehabil*. 2016;43(10):729–36 (**Epub 2016 Aug 30**).
50. Bendixen KH, Gjørup H, Baad-Hansen L, Dahl Hald J, Harsløf T, Schmidt MH, Langdahl BL, Haubek D. Temporomandibular disorders and psychosocial status in osteogenesis imperfecta: a cross-sectional study. *BMC Oral Health*. 2018;18(1):35.
51. Sawin PD, Menezes AH. Basilar invagination in osteogenesis imperfecta and related osteochondrodysplasias: medical and surgical management. *J Neurosurg*. 1997;86(6):950–60.
52. Arponen H, Mäkitie O, Haukka J, Ranta H, Ekholm M, Mäyränpää MK, Kaitila I, Waltimo-Sirén J. Prevalence and natural course of craniocervical junction anomalies during growth in patients with osteogenesis imperfecta. *J Bone Miner Res*. 2012;27(5):1142–9.
53. Arponen H, Vuorimies I, Haukka J, Valta H, Waltimo-Sirén J, Mäkitie O. Cranial base pathology in pediatric osteogenesis imperfecta patients treated with bisphosphonates. *J Neurosurg Pediatr*. 2015;15(3):313–20 (**Epub 2015 Jan 10**).
54. Reznikov N, Dagdeviren D, Tamimi F, Glorieux F, Rauch F, Retrouvey JM. Cone-beam computed tomography of osteogenesis imperfecta types III and IV: three-dimensional evaluation of craniofacial features and upper airways. *JBMR Plus*. 2019;3(6): e10124.
55. Simm PJ, Biggin A, Zacharin MR, Rodda CP, Tham E, Siafarikas A, Jefferies C, Hofman PL, Jensen DE, Woodhead H, Brown J, Wheeler BJ, Brookes D, Lafferty A, Munns CF, APEG Bone Mineral Working Group. Consensus guidelines on the use of bisphosphonate therapy in children and adolescents. *J Paediatr Child Health*. 2018;54(3):223–33.
56. Mueller B, Engelbert R, Baratta-Ziska F, Bartels B, Blanc N, Brizola E, Fraschini P, Hill C, Marr C, Mills L, Montpetit K, Pacey V, Molina MR, Schuurung M, Verhille C, de Vries O, Yeung EHK, Semler O. Consensus statement on physical rehabilitation in children and adolescents with osteogenesis imperfecta. *Orphanet J Rare Dis*. 2018;13(1):158.
57. Antoniazzi F, Mottes M, Fraschini P, et al. Osteogenesis Imperfecta. *Paediatr Drugs*. 2000;2:465–88.
58. Hald JD, Folkestad L, Swan CZ, Wanscher J, Schmidt M, Gjørup H, Haubek D, Leonhard CH, Larsen DA, Hjortdal JØ, Harsløf T, Duno M, Lund AM, Jensen JB, Brixen K, Langdahl B. Osteogenesis imperfecta and the teeth, eyes, and ears: a study of non-skeletal phenotypes in adults. *Osteoporos Int*. 2018;29(12):2781–9.
59. Ma MS, Najirad M, Taqi D, Retrouvey JM, Tamimi F, Dagdeviren D, Glorieux FH, Lee B, Sutton VR, Rauch F, Esfandiari S. Caries prevalence and experience in individuals with osteogenesis imperfecta: a cross-sectional multicenter study. *Spec Care Dentist*. 2019;39(2):214–9 (**Epub 2019 Feb 13**).
60. Malmgren B, Lindskog S. Assessment of dysplastic dentin in osteogenesis imperfecta and dentinogenesis imperfecta. *Acta Odontol Scand*. 2003;61(2):72–80.
61. Majorana A, Bardellini E, Brunelli PC, Lacaíta M, Cazzolla AP, Favia G. Dentinogenesis imperfecta in children with osteogenesis imperfecta: a clinical and ultrastructural study. *Int J Paediatr Dent*. 2010;20(2):112–8.
62. Johansson A, Omar R. Identification and management of tooth wear. *Int J Prosthodont*. 1994;7(6):506–16.
63. Loomans B, Opdam N, Attin T, Bartlett D, Edelhoff D, Frankenberger R, Benic G, Ramseyer S, Wetselaar P, Sterenberg B, Hickel R, Pallesen U, Mehta S, Banerji S, Lussi A, Wilson N. Severe tooth wear: European consensus statement on management guidelines. *J Adhes Dent*. 2017;19(2):111–9.
64. Ulseth JO, Hestnes A, Stovner LJ, Storhaug K. Dental caries and periodontitis in persons with Down syndrome. *Spec Care Dentist*. 1991;11(2):71–3.
65. Bandyopadhyay D. From mouth-level to tooth-level DMFS: conceptualizing a theoretical framework. *J Dent Oral Craniofac Epidemiol*. 2013;1(1):3–8.
66. Ainamo J, Barmes D, Beagrie G, Cutress T, Martin J, Sardo-Infirri J. Development of the World Health Organisation (WHO) community periodontal index of treatment needs (CPITN). *Int Dent J*. 1982;32:281–91.
67. Baelum V, Papapanou PN. CPITN and the epidemiology of periodontal disease. *Community Dent Oral Epidemiol*. 1996;24:367–8.
68. Bartlett D, Ganss C, Lussi A. Basic Erosive Wear Examination (BEWE): a new scoring system for scientific and clinical needs. *Clin Oral Investig*. 2008;12(Suppl 1):S65–8 (**Epub 2008 Jan 29**).
69. Smith BG, Knight JK. An index for measuring the wear of teeth. *Br Dent J*. 1984;156(12):435–8.
70. Van Wijk AJ, Tan SP. A numeric code for identifying patterns of human tooth agenesis: a new approach. *Eur J Oral Sci*. 2006;114(2):97–101.

71. de La Dure-Molla M, Fournier BP, Manzanares MC, Acevedo AC, Hennekam RC, Friedlander L, Boy-Lefèvre ML, Kerner S, Toupenay S, Garrec P, Vi-Fane B, Felizardo R, Berteretche MV, Jordan L, Ferré F, Clauss F, Jung S, de Chalendar M, Troester S, Kawczynski M, Chaloyard J; International Group of Dental Nomenclature; Manière MC, Berdal A, Bloch-Zupan A. Elements of morphology: standard terminology for the teeth and classifying genetic dental disorders. *Am J Med Genet A*. 2019;179(10):1913–1981. Epub 2019 Aug 29.
72. Raghoebar GM, Boering G, Vissink A, Stegenga B. Eruption disturbances of permanent molars: a review. *J Oral Pathol Med*. 1991;20(4):159–66.
73. Hennessy J, Al-Awadhi EA, Dwyer LO, Leith R. Treatment of ectopic first permanent molar teeth. *Dent Update*. 2012;39(9):656–8.
74. Verheij KL, Duggal M, Mejäre I, Papagiannoulis L, Koch G, Martens LC, Hal-lonsten AL. Judgement criteria for molar incisor hypomineralisation (MIH) in epidemiologic studies: a summary of the European meeting on MIH held in Athens, 2003. *Eur J Paediatr Dent*. 2003;4(3):110–3.
75. Kinaan BK. Overjet and overbite distribution and correlation: a comparative epidemiological English-Iraqi study. *Br J Orthod*. 1986;13(2):79–86.
76. Daniels C, Richmond S. The development of the index of complexity, outcome and need (ICON). *J Orthod*. 2000;27(2):149–62. <https://doi.org/10.1093/ortho/27.2.149>.
77. Brook PH, Shaw WC. The development of an index of orthodontic treatment priority. *Eur J Orthod*. 1989;11(3):309–20.
78. Schiffman E, Ohrbach R, Truelove E, Look J, Anderson G, Goulet JP, List T, Svensson P, Gonzalez Y, Lobbezoo F, Michelotti A, Brooks SL, Ceusters W, Drangsholt M, Ettlín D, Gaul C, Goldberg LJ, Haythornthwaite JA, Hollender L, Jensen R, John MT, De Laat A, de Leeuw R, Maixner W, van der Meulen M, Murray GM, Nixdorf DR, Palla S, Petersson A, Pionchon P, Smith B, Visscher CM, Zakrzewska J, Dworkin SF; International RDC/TMD Consortium Network, International association for Dental Research; Orofacial Pain Special Interest Group, International Association for the Study of Pain. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for Clinical and Research Applications: recommendations of the International RDC/TMD Consortium Network* and Orofacial Pain Special Interest Group†. *J Oral Facial Pain Headache*. 2014 Winter;28(1):6–27.
79. Rongo R, Ekberg E, Nilsson IM, Al-Khotani A, Alstergren P, Conti PCR, Durham J, Goulet JP, Hirsch C, Kalaykova SI, Kapos FP, Komiyama O, Koutris M, List T, Lobbezoo F, Ohrbach R, Peck CC, Restrepo C, Rodrigues MJ, Sharma S, Svensson P, Visscher CM, Wahlund K, Michelotti A. Diagnostic criteria for temporomandibular disorders (DC/TMD) for children and adolescents: an international Delphi study-Part 1-Development of Axis I. *J Oral Rehabil*. 2021;48(7):836–45 (**Epub 2021 May 19**).
80. Stegenga B, de Bont LG, de Leeuw R, Boering G. Assessment of mandibular function impairment associated with temporomandibular joint osteoarthritis and internal derangement. *J Orofac Pain*. 1993;7(2):183–95.
81. Broder HL, Wilson-Genderson M, Sischo L. Reliability and validity testing for the Child Oral Health Impact Profile-Reduced (COHIP-SF 19). *J Public Health Dent*. 2012;72(4):302–12. Epub 2012 Apr 27. Erratum in: *J Public Health Dent*. 2013 Winter;73(1):86.
82. Do LG, Spencer AJ. Evaluation of oral health-related quality of life questionnaires in a general child population. *Community Dent Health*. 2008;25(4):205–10.
83. Slade GD. Derivation and validation of a short-form oral health impact profile. *Community Dent Oral Epidemiol*. 1997;25(4):284–90.
84. Gilchrist F, Rodd H, Deery C, Marshman Z. Assessment of the quality of measures of child oral health-related quality of life. *BMC Oral Health*. 2014;23(14):40.
85. Larsson P, John MT, Nilner K, Bondemark L, List T. Development of an Orofacial Esthetic Scale in prosthodontic patients. *Int J Prosthodont*. 2010;23(3):249–56.
86. Larsson P, John MT, Nilner K, List T. Reliability and validity of the Orofacial Esthetic Scale in prosthodontic patients. *Int J Prosthodont*. 2010;23(3):257–62.
87. Porritt J, Morgan A, Rodd H, Gupta E, Gilchrist F, Baker S, Newton T, Creswell C, Williams C, Marshman Z. Development and evaluation of the children's experiences of dental anxiety measure. *Int J Paediatr Dent*. 2018;28(2):140–51 (**Epub 2017 Jul 29**).
88. Wong HM, Humphris GM, Lee GT. Preliminary validation and reliability of the Modified Child Dental Anxiety Scale. *Psychol Rep*. 1998;83(3 Pt 2):1179–86.
89. Howard KE, Freeman R. Reliability and validity of a faces version of the Modified Child Dental Anxiety Scale. *Int J Paediatr Dent*. 2007;17(4):281–8.
90. Al-Namankany A, de Souza M, Ashley P. Evidence-based dentistry: analysis of dental anxiety scales for children. *Br Dent J*. 2012;212(5):219–22.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.